

Topologically Linked Polymers are Anyon Systems

Franco Ferrari^{1,2,*}

¹*Institute of Physics, University of Szczecin, ul. Wielkopolska 15, 70-451 Szczecin, Poland*

²*Max Planck Institute for Polymer Research, 10 Ackermannweg, 55128 Mainz, Germany*

We consider the statistical mechanics of a system of topologically linked polymers, such as for instance a dense solution of polymer rings. If the possible topological states of the system are distinguished using the Gauss linking number as a topological invariant, the partition function of an ensemble of N closed polymers coincides with the $2N$ point function of a field theory containing a set of N complex replica fields and Abelian Chern-Simons fields. Thanks to this mapping to field theories, some quantitative predictions on the behavior of topologically entangled polymers have been obtained by exploiting perturbative techniques. In order to go beyond perturbation theory, a connection between polymers and anyons is established here. It is shown in this way that the topological forces which maintain two polymers in a given topological configuration have both attractive and repulsive components. When these opposite components reach a sort of equilibrium, the system finds itself in a self-dual point similar to that which, in the Landau-Ginzburg model for superconductors, corresponds to the transition from type I to type II superconductivity. The significance of self-duality in polymer physics is illustrated considering the example of the so-called $4 - plat$ configurations, which are of interest in the biochemistry of DNA processes like replication, transcription and recombination. The case of static vortex solutions of the Euler-Lagrange equations is discussed.

In this letter we consider the statistical mechanics of a system of topologically linked polymers[1], such as for instance a dense solution of polymer rings (catenanes). If the possible topological states of the system are distinguished using the Gauss linking number as a topological invariant, it has been shown in [2] that the partition function of an ensemble of N closed polymers coincides with the $2N$ point function of a field theory containing a set of N complex replica fields and Abelian Chern-Simons (C-S) fields. Thanks to this mapping to field theories, some quantitative predictions on the behavior of topologically entangled polymers have been obtained by exploiting perturbative methods, see Ref. [3] and references therein. If one wishes to go beyond perturbation theory, however, one is faced with the problem of field theories which contain multi-component scalar fields and gauge fields interacting together. Clearly, it is not easy to study theories of this kind even in the simplest case $N = 2$. The idea which we wish to pursue here in order to circumvent such difficulties is to establish a link between polymers and anyons [4]. Anyon field theories have been in fact intensively investigated. It is known for instance that their classical equations of motion admit vortex solutions [5]. In a special region of the space of parameters, called the Bogoml'nyi self-dual point [6], the attractive and repulsive forces between vortices vanish. This point corresponds in the Landau-Ginzburg model for superconductors to the transition from type I to type II superconductivity.

The first obstacle in this program is that anyons are living in $2 + 1$ dimensions, while polymer trajectories are intrinsically defined in three dimensions. For this reason, one coordinate must be singled out and regarded as *time*. Accordingly, a point of the three-dimensional Euclidean space \mathbf{R}^3 will be identified by a three vec-

tor $\xi^\mu = (\mathbf{r}, t)$, where $\mu = 1, 2, 3$, $\mathbf{r} = (x^1, x^2)$ and $x_3 = t$ plays the role of time. We need also to introduce the two-dimensional completely antisymmetric tensor ϵ^{ij} , $i = 1, 2$ and its three dimensional counterpart $\epsilon^{\mu\nu\rho}$, $\mu, \nu, \rho = 1, 2, 3$. These tensors are uniquely defined by the following conventions: $\epsilon^{12} = 1$ and $\epsilon^{123} = 1$. We use middle latin letters as two-dimensional space indices and middle greek letters as three dimensional indices. Sum over repeated space indices will be everywhere understood. The trajectories Γ_a of the polymers, $a = 1, \dots, N$, will be treated as continuous curves. Usually, these trajectories are parametrized by means of their arc-length σ_a : $\Gamma_a = \{(\mathbf{r}_a(\sigma_a), t_a(\sigma_a)) | 0 \leq \sigma_a \leq L_a\}$, L_a being the total length of the trajectory. To make a connection with anyons, however, it is convenient to use the time t as a parameter. Of course, the coordinate t is not a good parameter if we do not introduce a suitable of *sectioning* for the curves Γ_a . To this purpose, let us notice that, with respect to the t -direction, each curve Γ_a has maxima and minima defined by the condition: $\frac{dt_a(\sigma_a)}{d\sigma_a} = 0$. The number of minima, s_a , is equal to the number of maxima. Let us pick up a given point of minimum on Γ_a and call it $\tau_{a,1}$. Starting from $\tau_{a,1}$ and going along the curve after choosing arbitrarily a direction, one will encounter successively a point of maximum, that will be called $\tau_{a,2}$, a point of minimum $\tau_{a,3}$ and so on. In this way we obtain a set of $2s_a$ points $\tau_{a,1}, \dots, \tau_{a,2s_a}$. Now it is possible to split Γ_a into s_a open paths $\Gamma_{a,1}, \dots, \Gamma_{a,s_a}$ which connect pairwise contiguous points of maxima and minima:

$$\Gamma_{a,I_a} = \left\{ (\mathbf{r}_{a,I_a}(t), t) \left| \begin{array}{l} \tau_{a,I} \leq t \leq \tau_{a,I+1} \\ \mathbf{r}_{a,I_a}(\tau_{a,I_a+1}) = \mathbf{r}_{a,I_a+1}(\tau_{a,I_a+1}) \\ \mathbf{r}_{a,I_a}(\tau_{a,I_a}) = \mathbf{r}_{a,I_a-1}(\tau_{a,I_a}) \end{array} \right. \right\} \quad (1)$$

where I_a is a cyclic index such that $I_a \in \{1, \dots, 2s_a\}$ and $I_a + 2s_a = I_a$. The points $\mathbf{r}_{a,I_a}(\tau_{a,I_a})$ are considered as fixed points. This sectioning procedure is explicitly illustrated in the case $s = 3$ by Fig.; 1. By construction,




FIG. 1: Sectioning procedure for a $2s$ -plat Γ_a with $s = 3$.

there is a one-to-one correspondence between the points of the t -axis in the interval $[\tau_{a,I}, \tau_{a,I+1}]$ and the points of the trajectories $\Gamma_{a,I}$. As a consequence, it is possible to parametrize the trajectories Γ_{a,I_a} by means of the variable t .

We are now ready to construct the partition function of a system of topologically entangled polymers. For the sake of clarity, it is convenient to use some of the nomenclature of knot theory. In particular, from the mathematical point of view, each closed trajectory Γ_a can be regarded as a knot, while an ensemble of two or more trajectories defines a link. Moreover, with some slight abuse of language, a knot or a link with $2s$ points of maxima and minima will be called a $2s$ -plat. The concept of plats arises quite naturally in the processes of replication, recombination and transcription of DNA [8].

To start with, let us write the partition function for a link with the topology of a $2S$ -plat, where $S = \sum_{a=1}^n s_a$. The integers s_a are now kept fixed, so that the link is composed by knots Γ_a which are $2s_a$ -plats. In order to avoid a proliferation of indices, we will restrict ourselves to a system of two linked polymers, thus setting $N = 2$. Actually, this is not a big limitation, because the two-polymer problem contains all the ingredients of the N -polymer problem. Putting $\iota = \sqrt{-1}$, the desired polymer partition function is given by:

$$\mathcal{Z}(\lambda) = \int \mathcal{D}B_\mu \mathcal{D}C_\nu e^{-\iota S_{CS}} \int_{\text{boundary conditions}} \mathcal{D}\mathbf{r}_{a,I_a}(t) e^{-\mathcal{A}} \quad (2)$$

The boundary conditions on the trajectories Γ_{a,I_a} are those of Eq. (1). These boundary conditions are required by the fact that the open paths Γ_{a,I_a} , with a fixed and $I_a = 1, \dots, 2s_a$, must join together in a suitable way

at the points of maxima and minima τ_{a,I_a} in order to reconstruct the closed curve Γ_a .

The polymer action \mathcal{A} can be divided into two contributions:

$$\mathcal{A} = \mathcal{A}_{free} + \mathcal{A}_{top} \quad (3)$$

The free part \mathcal{A}_{free} is:

$$\mathcal{A}_{free} = \sum_{a=1}^2 \sum_{I_a=1}^{2s_a} \int_{\tau_{a,I_a}}^{\tau_{a,I_a+1}} dt (-1)^{I_a-1} g_{a,I_a} \left| \frac{d\mathbf{r}_{a,I_a}(t)}{dt} \right| \quad (4)$$

The factors $(-1)^{I_a-1}$ appearing in Eq. (4) are necessary to make \mathcal{A}_{free} positive definite. The parameters g_{a,I_a} are related to the lengths L_{a,I_a} of the open chains Γ_{a,I_a} as follows. Let us discretize Γ_{a,I_a} considering it as a random chains with n segments. If Eq. (4) would be the action of a set of free two-dimensional chains, in the limit of large values of n and keeping fixed the ratio $n/g_{a,I_a}$, one would obtain the well known result [7]: $L_{a,I_a} \sim n/g_{a,I_a}$. Here things are a little bit more complicated, because one has to take into account the third dimension, which in turn is also the variable which parametrizes the curves. As a consequence, the above relation governing the length of the trajectories is replaced by:

$$L_{a,I_a}^2 \sim |\tau_{a,I_a+1} - \tau_{a,I_a}|^2 + \frac{n}{g_{a,I_a}} |\tau_{a,I_a+1} - \tau_{a,I_a}| \quad (5)$$

The second term in Eq. (3) is:

$$\begin{aligned} \mathcal{A}_{top} = & \iota \lambda \sum_{I=1}^{2s_1} \int_{\tau_{1,I}}^{\tau_{1,I+1}} dt \frac{d\xi_{1,I}^\mu(t)}{dt} B_\mu(\mathbf{r}_{1,I}(t), t) \\ & + \iota \frac{\kappa}{2\pi} \sum_{J=1}^{2s_2} dt \frac{d\xi_{2,J}^\mu(t)}{dt} C_\mu(\mathbf{r}_{2,J}(t), t) \end{aligned} \quad (6)$$

In the above equation $B_\mu = (\mathbf{B}, B_0)$ and $C_\mu = (\mathbf{C}, C_0)$ are two Abelian C-S vector fields. We have put $B_3 \equiv B_0$ and $C_3 \equiv C_0$ in order to stress the role of time of the third spatial coordinate. κ and λ are real parameters. κ coincides with the C-S coupling constant (see below). The spurious gauge degrees of freedom have been eliminated using the gauge condition (Coulomb gauge):

$$\nabla \cdot \mathbf{B} = \nabla \cdot \mathbf{C} = 0 \quad (7)$$

so that the gauge fixed C-S action S_{CS} looks as follows:

$$S_{CS} = \frac{\kappa}{4\pi} \int d^3\xi [B_0 \epsilon^{ij} \partial_i C_j + C_0 \epsilon^{ij} \partial_i B_j] \quad (8)$$

The C-S fields act as auxiliary fields which impose topological constraints on the closed trajectories Γ_1 and Γ_2 . To show this, one has to integrate out these fields from

the partition function (2) and then to perform an inverse Fourier transformation of $\mathcal{Z}(\lambda)$:

$$\mathcal{Z}(m) = \int_0^{2\pi} \frac{d\lambda}{2\pi} e^{im\lambda} \mathcal{Z}(\lambda) \quad (9)$$

The details of this calculation have been already explained in Ref. [3] and will not be repeated here. The result is the new partition function $\mathcal{Z}(m)$, which describes the fluctuations of two polymer rings constrained by the condition:

$$m = \chi(\Gamma_1, \Gamma_2) \quad (10)$$

where $\chi(\Gamma_1, \Gamma_2)$ is the Gauss linking number and the topological number m must be an integer. A few comments are in order at this point. First of all, in [3] the topological constraint (10) has been derived from the C-S path integral using the Lorentz gauge, while here the Coulomb gauge (7) has been chosen. Formally, things works exactly in the same way in both gauges. However, one should keep in mind that the final expression of the Gauss linking number $\chi(\Gamma_1, \Gamma_2)$ obtained starting from the Lorentz gauge differs from the expression that one obtains starting from the Coulomb gauge. Of course, the two expressions are equivalent due to gauge invariance, but the way in which the winding of one trajectory around the other is counted, is realized in a very different way. Another important point is that our approach allows a more refined classification of links than the bare Gauss linking number. Indeed, two links which have the same Gauss linking number can be further distinguished by their plat-structure, i. e. by the number of maxima and minima of the knots which are composing the links. Finally, it is not difficult to add also the repulsive steric interactions between the monomers in the partition function (2). It is interesting to notice that, since we have chosen the coordinate $x_3 \equiv t$ in order to parametrize the curves, the usual δ -function potential which describes these forces takes the form: $V(\mathbf{r}, t) = v_0 \delta(\mathbf{r}) \delta(t)$, v_0 being a temperature dependent constant. Due to the presence of the δ -function in time, the steric interactions become instantaneous in our case. In the following we will not

discuss the steric interactions, in part because we prefer to concentrate on the pure topological interactions, in part because the inclusion of these repulsive forces complicates enormously the task of finding non-perturbative analytic solutions of the two-polymer problem under consideration. Thus, we will assume that in our system steric interactions are negligible. This approximation is valid for instance in polymer solutions which are very dilute or in which the temperature is near the so-called theta-point. The hypothesis of very low monomer densities is natural in the present context, because we are discussing the statistical properties of two isolated polymer rings, which are supposed to be very long and thin.

To establish the connection with anyons, we need to pass in the partition function (2) from the statistical sum over the trajectories $\mathbf{r}_{a,I_a}(t)$ to a path integral over fields. The details of the derivation of the partition function $\mathcal{Z}(\lambda)$ in terms of fields in the general case and including also the steric interactions will be presented elsewhere. Here we will restrict ourselves to links which have the structure of 4-plats, putting henceforth $s_1 = s_2 = 1$. This case is particularly interesting for biological applications, since most of the links produced by in vitro enzymology experiments are 4-plats [8]. Moreover, "small" links up to seven crossings are all 4-plats.

Introducing a set of n_r complex replica fields:

$$\begin{aligned} \vec{\Psi}_{a,I_a} &= (\psi_{a,I_a}^1, \dots, \psi_{a,I_a}^{n_r}) \\ \vec{\Psi}_{a,I_a}^* &= (\psi_{a,I_a}^{1*}, \dots, \psi_{a,I_a}^{n_r*}) \end{aligned} \quad (11)$$

and a convenient notation for covariant derivatives:

$$\mathbf{D}(\pm\lambda, \mathbf{B}) = \nabla \pm i\lambda \mathbf{B} \quad \mathbf{D}(\pm\frac{\kappa}{2\pi}, \mathbf{C}) = \nabla \pm i\frac{\kappa}{2\pi} \mathbf{C} \quad (12)$$

the 4-plat partition function can be expressed as follows:

$$\mathcal{Z}_{4-plat}(\lambda) = \lim_{n \rightarrow 0} \int \mathcal{D}(fields) \prod_{I=1}^2 \mathcal{O}_{1,I} \prod_{J=1}^2 \mathcal{O}_{2,J} e^{-\mathcal{S}_{4-plat}} \quad (13)$$

The action \mathcal{S}_{4-plat} contains the free field action and the topological terms:

$$\begin{aligned} \mathcal{S}_{4-plat} &= \int_{\tau_{1,1}}^{\tau_{1,2}} dt \int d\mathbf{x} \left\{ \vec{\Psi}_{1,1}^* \partial_t \vec{\Psi}_{1,1} + \frac{1}{4g_{11}} \left| \mathbf{D}(-\lambda, \mathbf{B}) \vec{\Psi}_{1,1} \right|^2 + \vec{\Psi}_{1,2}^* \partial_t \vec{\Psi}_{1,2} + \frac{1}{4g_{12}} \left| \mathbf{D}(\lambda, \mathbf{B}) \vec{\Psi}_{1,2} \right|^2 \right\} + \\ &+ \int_{\tau_{2,1}}^{\tau_{2,2}} dt \int d\mathbf{x} \left\{ \vec{\Psi}_{2,1}^* \partial_t \vec{\Psi}_{2,1} + \frac{1}{4g_{21}} \left| \mathbf{D}(-\frac{\kappa}{2\pi}, \mathbf{C}) \vec{\Psi}_{2,1} \right|^2 + \vec{\Psi}_{2,2}^* \partial_t \vec{\Psi}_{2,2} + \frac{1}{4g_{22}} \left| \mathbf{D}(\frac{\kappa}{2\pi}, \mathbf{C}) \vec{\Psi}_{2,2} \right|^2 \right\} \end{aligned} \quad (14)$$

$d\mathbf{x} \equiv dx_1 dx_2$ is the infinitesimal volume element in the

two-dimensional space and $\partial_t = \frac{\partial}{\partial t}$. Let us note that in

the partition function (13) we have already integrated out the time components of the Chern-Simons fields B_0, C_0 , which play the role of Lagrange multipliers and impose the following constraints:

$$\mathcal{B} = 2 \left(\left| \vec{\Psi}_{2,1} \right|^2 - \left| \vec{\Psi}_{2,2} \right|^2 \right) \theta(\tau_{2,2} - t) \theta(t - \tau_{2,1}) \quad (15)$$

$$\mathcal{C} = \frac{4\pi}{\kappa} \left(\left| \vec{\Psi}_{1,1} \right|^2 - \left| \vec{\Psi}_{1,2} \right|^2 \right) \theta(\tau_{1,2} - t) \theta(t - \tau_{1,1}) \quad (16)$$

In the above equation \mathcal{B} and \mathcal{C} are the magnetic fields associated to the vector potentials \mathbf{B} and \mathbf{C} respectively:

$$\mathcal{B} = \partial_1 B_2 - \partial_2 B_1 = \epsilon^{ij} \partial_i B_j \quad (17)$$

$$\mathcal{C} = \partial_1 C_2 - \partial_2 C_1 = \epsilon^{ij} \partial_i C_j \quad (18)$$

and $\theta(t)$ is the Heaviside theta function $\theta(t) = 0$ if $t < 0$ and $\theta(t) = 1$ if $t \geq 0$. The symbol $\mathcal{D}(\text{fields})$ in Eq. (13) is a shorthand for the field integration measure:

$$\mathcal{D}(\text{fields}) = \int \prod_{I_a=1}^{2s_a} \mathcal{D}\vec{\Psi}_{a,I_a}^* \mathcal{D}\vec{\Psi}_{a,I_a} \quad (19)$$

Finally, the operators \mathcal{O}_{a,I_a} are given by:

$$\begin{aligned} \mathcal{O}_{a,I_a} &= \psi_{a,I_a}^1(\mathbf{r}_{a,I_a}(\tau_{a,I_a+1}), \tau_{a,I_a+1}) \\ &\quad \psi_{a,I_a}^{1*}(\mathbf{r}_{a,I_a}(\tau_{a,I_a}), \tau_{a,I_a}) \end{aligned} \quad (20)$$

Eqs. (13)–(20) establish the desired connection between polymers and anyons. In fact, the action (14), together with the constraints (15)–(16), coincides formally with the action of a multi-layered anyon system. The number of layers is $2(s_1 + s_2) = 4$, while the fields $\vec{\Psi}_{a,I_a}^*, \vec{\Psi}_{a,I_a}$

propagate anyonic particles with spin n_r and the statistics of boson. One difference from anyonic field theories is due to the fact that, in the case of real particles, one considers the evolution of the whole system from an initial time T_1 and a final time T_2 . Here, instead, time intervals measure the extensions of the trajectories Γ_{a,I_a} in the x_3 direction. For this reason, in the partition function (13) part of the system evolves within the time interval $[\tau_{1,1}, \tau_{1,2}]$ and another part within the time interval $[\tau_{2,1}, \tau_{2,2}]$. This inhomogeneity in the time intervals and the use of the Coulomb gauge, which makes the interactions instantaneous, is the cause of the presence of the Heaviside θ -functions in Eqs. (15)–(16). In this sense, the complete analogy with anyons can be recovered only in the limit

$$\tau_{1,1} = \tau_{2,1} \equiv T_1 \quad \tau_{1,2} = \tau_{2,2} = T_2 \quad (21)$$

Throughout the rest of this letter we will assume that the above condition is satisfied. This is to avoid technical complications in dealing with the θ -functions.

The connection to the anyon problem suggest to apply to the action of Eq. (14) the Bogomol'nyi transformations [6]. Using these transformation in a suitable way, it is possible to rewrite \mathcal{S}_{4-plat} in the following form:

$$\mathcal{S}_{4-plat} = \mathcal{F}_{4-plat} + \mathcal{I}_{4-plat} \quad (22)$$

Neglecting surface terms and introducing the notation $D_{\pm} = D_1 \pm iD_2$, where D_i , $i = 1, 2$, denotes the i -th space components of the covariant derivatives (12), one has that:

$$\begin{aligned} \mathcal{F}_{4-plat} &= \int_{T_1}^{T_2} dt \int d\mathbf{x} \left[\sum_{a=1}^2 \sum_{I_a=1}^2 \vec{\Psi}_{a,I_a}^* \partial_t \vec{\Psi}_{a,I_a} + \right. \\ &\quad \left. + \frac{1}{4g_{11}} \left| D_+(-\lambda, \mathbf{B}) \vec{\Psi}_{1,1} \right|^2 + \frac{1}{4g_{12}} \left| D_+(\lambda, \mathbf{B}) \vec{\Psi}_{1,2} \right|^2 + \frac{1}{4g_{21}} \left| D_-\left(-\frac{\kappa}{2\pi}, \mathbf{C}\right) \vec{\Psi}_{2,1} \right|^2 + \frac{1}{4g_{22}} \left| D_-\left(\frac{\kappa}{2\pi}, \mathbf{C}\right) \vec{\Psi}_{2,2} \right|^2 \right] \end{aligned} \quad (23)$$

and

$$\mathcal{I}_{4-plat} = \frac{\lambda}{2} \int_{T_1}^{T_2} dt \int d\mathbf{x} \left[\left(\frac{|\vec{\Psi}_{1,1}|^2}{g_{11}} - \frac{|\vec{\Psi}_{1,2}|^2}{g_{12}} \right) \times \left(\left| \vec{\Psi}_{2,1} \right|^2 - \left| \vec{\Psi}_{2,2} \right|^2 \right) - \left(\frac{|\vec{\Psi}_{2,1}|^2}{g_{12}} - \frac{|\vec{\Psi}_{2,2}|^2}{g_{22}} \right) \times \left(\left| \vec{\Psi}_{1,1} \right|^2 - \left| \vec{\Psi}_{1,2} \right|^2 \right) \right] \quad (24)$$

It is easy to recognize from Eq. (23) that \mathcal{F}_{4-plat} consists in the sum of the self-dual actions of four different families of anyons $\vec{\Psi}_{a,I_a}$. These families are coupled together by the constraints (15)–(16). The second contribution to \mathcal{S}_{4-plat} , the interaction term \mathcal{I}_{4-plat} , contains

instead Coulomb-like potentials, which attract or repel the monomers belonging to different trajectories Γ_{a,I_a} . It is difficult to guess what is the influence of these Coulomb forces on the statistical behavior of the polymers. The problem is that their strength is proportional to λ (see

Eq. (24)) and λ is not a real coupling constant with a physical meaning. At the end, we are interested to eliminate λ and to express the polymer partition function in terms of the topological number m , as we did in Eq. (9). m is in fact the true physical parameter which describes the topological state of the system. What is possible to conclude from Eqs. (22)–(24) is that topological forces have attractive and repulsive components, which interfere with the steric interactions. This result confirms at a non-perturbative level a previous result obtained using perturbative methods [3]. Also in Ref. [9] it has been shown in the limit in which fluctuations in the particle densities are relatively small that the C-S fields generate effective Coulomb interactions in anyon models.

Remarkably, our approach reveals that the two-polymer system under consideration has a self-dual point. This point is reached when the parameters g_{aI_a} are equal:

$$g_{11} = g_{12} = g_{21} = g_{22} = g \quad (25)$$

If the above relations are satisfied, it is easy to see that $\mathcal{I}_{4-plat} = 0$, so that the only surviving terms in the action \mathcal{S}_{4-plat} are those coming from the self-dual contributions in \mathcal{F}_{4-plat} . We note that the existence of the self-dual point (25) is not submitted to the assumption of Eq. (21). Indeed, let us suppose that the points of maxima and minima τ_{a,I_a} of the trajectories Γ_{a,I_a} are not aligned as specified by Eq. (21). The only difference in the expression of the interaction term \mathcal{I}_{4-plat} is that now the values of T_1 and T_2 , which give the boundaries of the integration over time, should be replaced by the following ones:

$$T_1 = \max[\tau_{1,1}, \tau_{2,1}] \quad T_2 = \min[\tau_{1,2}, \tau_{2,2}] \quad (26)$$

The choice (26) of integration limits in \mathcal{I}_{4-plat} is dictated by the presence of the θ -functions of Heaviside in the constraints (15)–(16). Physically, Eq. (26) is motivated by the fact that, in the Coulomb gauge, all interactions become instantaneous. Therefore, if maxima and minima are not aligned, it is easy to see that the trajectories may only interact in the time interval $[T_1, T_2]$, where T_1 and T_2 are given by Eq. (26). Substituting the new integration limits (26) in Eq. (24), it is clear that, if Eq. (25) is satisfied, the effective Coulomb interactions in \mathcal{I}_{4-plat} vanish identically, so that the action \mathcal{S}_{4-plat} coincides once again with the pure self-dual part \mathcal{F}_{4-plat} . Thus, to have self-duality we only need that the effects of the steric interactions are negligible, such as for instance in the case of very low monomer concentration or at the theta-point.

The self-duality condition (25) may be physically interpreted as the requirement that the trajectories Γ_{a,I_a} must be homogeneous, in the sense that they should have equal persistence lengths. In this case, the attractive and repulsive forces described by the term \mathcal{I}_{4-plat} counterbalance themselves in a sort of equilibrium which establishes

the self-dual regime. This does not mean however that attractions and repulsions between monomers disappear, because Coulomb-like forces still hide in the self-dual part of the action \mathcal{F}_{4-plat} .

We would like now to exploit further the analogy of the polymer problem with anyons field theories and search for possible non-trivial classical solutions which minimize the polymer free energy \mathcal{S}_{4-plat} . Of course, one should keep in mind that, even in the case of the simplest model of anyons, vortex solutions outside the self-dual point have been found only by means of numerical methods. Also at the self-dual point, there is no clear and detailed understanding of the dynamics of C-S vortices [10]. Here the situation is further complicated by the presence of replica fields and multiple families of anyons which are mixed together due to the constraints (15)–(16). In view of these limitations, it seems reasonable to restrict ourselves to the self-duality regime (25) and to static solutions, which satisfy the conditions $\partial_t \psi_{a,I_a}^\omega = 0$ for every value of $a = 1, 2$, $I = 1, 2$ and $\omega = 1, \dots, n_r$. As a consequence, from now on we will consider the pure self-dual action:

$$\begin{aligned} \mathcal{S}_{4-plat} = & \frac{(T_2 - T_1)}{4g} \int d\mathbf{x} \left[\left| D_+(-\lambda, \mathbf{B}) \vec{\Psi}_{1,1} \right|^2 + \left| D_+(\lambda, \mathbf{B}) \vec{\Psi}_{1,2} \right|^2 + \right. \\ & \left. \left| D_-\left(-\frac{\kappa}{2\pi}, \mathbf{C}\right) \vec{\Psi}_{2,1} \right|^2 + \left| D_-\left(\frac{\kappa}{2\pi}, \mathbf{C}\right) \vec{\Psi}_{2,2} \right|^2 \right] \quad (27) \end{aligned}$$

Of course, static solutions do not fit very well with the physical boundary conditions which should be imposed to the classical fields at the points of maxima and minima T_1 and T_2 . For this reason, we are supposing here implicitly that polymers are very long and that the interval $T_2 - T_1$ is large. In this way the relevant contributions to the free energy \mathcal{S}_{4-plat} come from instants t which are far enough from the extrema located at $t = T_1$ and $t = T_2$. At these intermediate points the trajectories fluctuate in such a way that, at each fixed instant t , they visit more or less with the same frequency the same locations of the two dimensional space spanned by the vectors $\mathbf{r}_{a,I_a}(t)$.

From the action (27) one finds the following Euler-Lagrange equations of motion:

$$D_+(-\lambda, \mathbf{B}) \psi_{1,1}^\omega = D_+(\lambda, \mathbf{B}) \psi_{1,2}^\omega = 0 \quad (28)$$

$$D_-\left(-\frac{\kappa}{2\pi}, \mathbf{C}\right) \psi_{2,1}^\omega = D_-\left(\frac{\kappa}{2\pi}, \mathbf{C}\right) \psi_{2,2}^\omega = 0 \quad (29)$$

To these equations one should also add the constraints (15)–(16) which determine the transverse components of the fields \mathbf{B}, \mathbf{C} and the Coulomb gauge fixing (7). In total, after separating the real and imaginary parts in Eqs. (28)–(29), we obtain $12 \times n_r$ equations, which completely specify the classical field configurations. It seems natural to try replica symmetric solutions of the kind:

$$\psi_{a,I_a}^\omega = \psi_{a,I_a} \quad \psi_{a,I_a}^{\omega*} = \psi_{a,I_a}^* \quad (30)$$

where $a = 1, 2$, $I_a = 1, 2$ and $\omega = 1, \dots, n_r$. Moreover, we switch to polar coordinates in order to express the complex fields:

$$\psi_{a,I_a} = e^{i\phi_{a,I_a}} \rho_{a,I_a}^{1/2} \quad (31)$$

The consistency of Eqs. (28)–(29) requires that:

$$\phi_{a,1} = -\phi_{a,2} \quad \rho_{a,1} = \frac{c_a}{\rho_{a,2}} \quad (32)$$

Here c_1 and c_2 are arbitrary constants different from zero. Also the densities ρ_{a,I_a} must not vanish. There are other ways to make Eqs. (28)–(29) consistent, but they lead either to unphysical or to trivial solutions. After imposing the consistency conditions (32) in Eqs. (28)–(29), there remain still four independent relations, which can be used to fix the components of the vector fields \mathbf{B} , \mathbf{C} :

$$\lambda B_i = \partial_i \phi_{1,1} + \frac{1}{2} \epsilon_{ij} \partial^j \ln \rho_{1,1} \quad (33)$$

$$\frac{\kappa}{2\pi} C_i = \partial_i \phi_{2,1} - \frac{1}{2} \epsilon_{ij} \partial^j \ln \rho_{2,1} \quad (34)$$

It is easy to prove that the Coulomb gauge requirement (7) is satisfied only if the phases $\phi_{1,1}$ and $\phi_{2,1}$ are constant. Finally, inserting Eqs. (33)–(34) in the constraints (15)–(16), it is possible to determine the remaining degrees of freedom $\rho_{1,1}$ and $\rho_{2,1}$:

$$\Delta \ln \rho_{1,1} = 4\lambda n_r \left(\frac{c_2}{\rho_{2,1}} - \rho_{2,1} \right) \quad (35)$$

$$\Delta \ln \rho_{2,1} = 4\lambda n_r \left(\rho_{1,1} - \frac{c_1}{\rho_{1,1}} \right) \quad (36)$$

A further study of the above two equations requires numerical methods.

Concluding, we have succeeded in establishing a connection between topologically linked polymers and anyon field theories. We have limited ourselves to 4-plats, but it is possible to generalize our results to the case of any $2s$ -plat. In our approach, links which belong to different $2s$ -plat configurations are distinguished. This allows a better classification of the topological states of polymers than the bare Gauss linking number. In view of possible applications to knot theory, it would be nice to extend also to non-abelian C-S field theories our techniques based on the Coulomb gauge and on the sectioning procedure of loops described above.

The analogy between polymers and anyons has been exploited to investigate the statistical mechanics of 4-plats, a particular class of links, which is relevant in biological applications. It turns out that, as anyon particles in multi-layered systems, also 4-plats have a self-dual point. Within the hypothesis of Eq. (21), which demands that the points of minima $\tau_{a,2I_a+1}$ are aligned at the same height $t = T_1$, while the points of maxima

are aligned at the height $t = T_2$, self-duality occurs when all open curves Γ_{a,I_a} have the same length. Particular polymer configurations of this kind may exist in nature, for instance after the process of DNA replication. However, we have seen that the requirements for the presence of a self-dual point are much more general: We only need that the effects of the steric interactions are negligible, such as for instance in the case of very low monomer concentration or at the theta-point.

The existence of the self-dual point (25) shows that the physics of interacting polymers is much richer than previously expected. For instance, the classical equations of motion admit non-trivial solutions, which are the analogs of vortex-like solutions in anyon models. By analytical methods it is just possible to investigate static field configurations. These suffer the limitations mentioned after Eq. (27) and cannot be defined if the requirements of Eqs. (21) are not fulfilled. In fact, if maxima and minima are localized at different heights on the t -axis, the action \mathcal{S}_{4-plat} becomes explicitly dependent on time due to the presence of the Heaviside θ -functions in the fields \mathbf{B} and \mathbf{C} . This time dependence is very mild. Its only effect is that the boundaries of integration over time in the various terms composing the action \mathcal{S}_{4-plat} do not coincide. Yet, this is sufficient to spoil the possibility of having purely static field configurations. For these reasons, any conclusion on the meaning of the static solutions in the case of polymers should be taken with some care. Having in mind these caveats, it is interesting to note that, in the solutions which we have studied here, it seems to prevail a repulsive force between couples of trajectories Γ_{a,I_a} belonging to the same polymer. This is what Eq. (32) suggests, because the densities $\rho_{a,1}$ are constrained to be inversely proportional to their counterparts $\rho_{a,2}$ at each point \mathbf{x} .

It is worthing to point out that, despite their close resemblances, the mechanisms used to establish the self-dual regime in our polymer model and in usual anyon field theories are different. In anyon field theories, in fact, the existence of the self-dual point requires that the effective Coulomb interactions, which arise after performing the Bogomol'nyi transformations, are canceled against true Coulomb interactions of charged particles [11]. In the polymer model, instead, the effective Coulomb interactions, represented by the term \mathcal{I}_{4-plat} , counterbalance themselves and the presence of other interactions is not needed. This self-balancing transition to self-duality may occur also in other physical problems where multi-layered systems of non-charged particles are relevant. Another application of our approach are polymer brushes. In that case, C-S field theories in the Coulomb gauge are able to take into account the effects related to the winding of neighboring polymer trajectories.

Part of this work has been carried out during a visit to the Max Planck Institute for Polymer Research. This visit has been funded by a grant from the German Aca-

demic Exchange Service (DAAD), which is gratefully acknowledged. The author wishes also to thank the Max Planck Institute for Polymer Research for the nice hospitality. He is also indebted to V. A. Rostiashvili and T. Vilgis, which participated in the early stages of this work, for their invaluable help and their constant encouragement.

* Electronic address: ferrari@univ.szczecin.pl

- [1] For a review on topologically linked polymers see for instance:
A. L. Kholodenko and T. A. Vilgis, *Phys. Rep.* **298** (1998), 251.
- [2] F. Ferrari, H. Kleinert and I. Lazzizzera, *Int. J. Mod. Phys. B* **14** (2000) 3881 [arXiv:cond-mat/0005300].
- [3] F. Ferrari, *Annalen Phys.* **11** (2002) 255.
- [4] F. Wilczek, *Phys. Rev. Lett.* **49**, 957 (1982); F. Wilczek and A. Zee, *ibid.* **51** (1983), 2250.
- [5] R. Jackiw and S.-Y. Pi, *Phys. Rev.* **D42** (1990), 3500.
- [6] E. Bogomol'nyi, *Sov. J. Nucl. Phys.* **24** (1976), 449.
- [7] H. Kleinert, *Path Integrals in Quantum Mechanics, Statistics, Polymer Physics, and Financial Markets*, (World Scientific Publishing, 3rd Ed., Singapore, 2003).
- [8] S. A. Wasserman and N. R. Cozzarelli, *Science* **232** (1986), 952; D. W. Sumners, *Notices of the Am. Math. Soc.* **42** (5) (1995), 528.
- [9] Y.-H. Chen, F. Wilczek and E. Witten, *Int. Jour. Mod. Phys. B* **3** (7) (1989), 1001.
- [10] G. Dunne, *Aspect of Chern-Simons Theories*, published in *Topological Aspects of Low Dimensional Systems*, A. Comtet et al. (Eds), Springer Verlag 2000.
- [11] Z. F. Ezawa and A. Iwazaki, *Phys. Rev.* **B47** (1993), 7295; S. M. Girvin and A. H. MacDonald, *Perspectives in Quantum Hall Effects*, S. D. Sarma and A. Pinczuk (Eds), Chap. 5, Wiley, New York, 1997.